arXiv:1412.5817v2 [math.AT] 20 Jun 2015

Fixed point indices of central configurations

D.L. Ferrario

June 23, 2015

To Professor Andrzej Granas

Abstract

Central configurations of n point particles in $E \approx \mathbb{R}^d$ with respect to a potential function U are shown to be the same as the fixed points of the normalized gradient map $F = -\nabla_M U/\|\nabla_M U\|_M$, which is an $SO(d)$-equivariant self-map defined on the inertia ellipsoid. We show that the $SO(d)$-orbits of fixed points of F are all fixed points of the map induced on the quotient by $SO(d)$, and give a formula relating their indices (as fixed points) with their Morse indices (as critical points). At the end, we give an example of a non-planar relative equilibrium which is not a central configuration.
MSC Subject Class: 55M20, 37C25, 70F10.
Keywords: Central configurations, relative equilibria, n-body problem.

1 Central configurations

Let $E = \mathbb{R}^d$ be the d-dimensional Euclidean space, and $X = E^n \smallsetminus \Delta$ the *configuration space* of E, defined as the space of all n-tuples of distinct points $\boldsymbol{q} = (\boldsymbol{q}_1, \dots, \boldsymbol{q}_n) \in E^n$ such that $\boldsymbol{q} \notin \Delta$, where Δ is the collision set

$$\Delta = \bigcup_{i<j} \{\boldsymbol{q} \in E^n : \boldsymbol{q}_i = \boldsymbol{q}_j\} \ .$$

Let $U\colon X \to \mathbb{R}$ be a regular potential function. For example, the gravitational potential

$$U(\boldsymbol{q}) = \sum_{i<j} \frac{m_i m_j}{\|\boldsymbol{q}_i - \boldsymbol{q}_j\|^\alpha}, \tag{1.1}$$

or more general potentials (charged particles, ...).

Given n real non-zero numbers $m_1, \dots, m_n$ (representing the *masses* of the n interacting point particles), let $\langle -, - \rangle_M$ denote the *mass-metric* on E^n, defined as $\langle \boldsymbol{v}, \boldsymbol{w} \rangle_M = \sum_{j=1}^n m_j \boldsymbol{v}_j \cdot \boldsymbol{w}_j$, where $\boldsymbol{v}, \boldsymbol{w} \in \mathbb{R}^{dn}$ are two vectors tangent to E^n, and $\cdot$ denotes the standard scalar product in E. If the

masses are positive, then the mass-metric is a non-degenerate scalar product on E^n, which yields both the kinetic quadratic form (on the tangent bundle) $2K = \|\dot{\boldsymbol{q}}\|_M^2 = \sum_i m_i \|\dot{\boldsymbol{q}}_i\|^2$ and the inertia form $I = \|\boldsymbol{q}\|_M^2 = \sum_i m_i \|\boldsymbol{q}_i\|^2$.

Given such m_i, let ∇_M denote the gradient of U with respect to the bilinear product $\langle -, - \rangle_M$. Let us recall that if $dU(\boldsymbol{q})$ denotes the differential of U evaluated in $\boldsymbol{q}$, then for each tangent vector $\boldsymbol{v} \in T_{\boldsymbol{q}} E^n$, $dU(\boldsymbol{q})[\boldsymbol{v}] = \langle \nabla_M U(\boldsymbol{q}), \boldsymbol{v} \rangle_M$, and therefore

$$dU(\boldsymbol{q})[\boldsymbol{v}] = \sum_j \frac{\partial U}{\partial \boldsymbol{q}_j} \cdot \boldsymbol{v}_j = \sum_j m_j \left(m_j^{-1} \frac{\partial U}{\partial \boldsymbol{q}_j} \right) \cdot \boldsymbol{v}_j,$$

from which it follows that in standard coordinates $(\nabla_M U)_j = m_j^{-1} \dfrac{\partial U}{\partial \boldsymbol{q}_j}$.

Given the mass-metric gradient $\nabla_M U$, the corresponding *Newton equations* are

$$\frac{d^2 \boldsymbol{q}}{dt^2} = \nabla_M U(\boldsymbol{q}). \tag{1.2}$$

(1.3) Definition. A configuration $\boldsymbol{q} \in X$ is a *central configuration* iff there exists $\lambda \in \mathbb{R}$, $\lambda \neq 0$, such that $\nabla_M U(\boldsymbol{q}) = \lambda \boldsymbol{q}$.

(1.4) Remark. If U is homogeneous, this is equivalent to: $\boldsymbol{q}$ is a central configuration iff there is a real-valued function $\varphi(t)$ such that $\varphi(t)\boldsymbol{q}$ is a solution of (1.2)[1]. If U is homogeneous, the set of central configurations is a cone in X.

(1.5) Remark. Furthermore, if U is invariant with respect to the group of all translations of E, then central configurations belong to the subspace

$$Y = \{\boldsymbol{q} \in X : \sum_j m_j \boldsymbol{q}_j = 0\} \subset X \subset E^n,$$

and $\forall \boldsymbol{q} \in X \implies \nabla_M U(\boldsymbol{q}) \in \overline{Y} \subset E^n$, where $\overline{Y}$ is the closure of Y in E^n. Therefore, if U is translation-invariant, the set of central configurations is a subset of Y. Sometimes central configurations are defined with the equation $\nabla_M U(\boldsymbol{q}) = \lambda(\boldsymbol{q} - \boldsymbol{c})$, where $\boldsymbol{c}$ is the center of mass $\boldsymbol{c} = (\sum_j m_j)^{-1} \sum_j m_j \boldsymbol{q}_j$ of the configuration $\boldsymbol{q}$. This equation is invariant with respect to translations (if U is so).

(1.6) Definition. A configuration $\boldsymbol{q} \in X$ is a *relative equilibrium* iff there is a one-parameter group of rotations $\varphi^t \colon E \to E$ (around the origin, without loss of generality) such that

$$\varphi^t(\boldsymbol{q}) = (\varphi^t(\boldsymbol{q}_1), \varphi^t(\boldsymbol{q}_2), \ldots, \varphi^t(\boldsymbol{q}_n))$$

satisfies the equations of motion (1.2).[2]

[1] Cf. [8], p. 61.
[2] Cf. [17], p. 47.

One-parameter subgroups of $SO(N)$ are of the form $\varphi^t(\boldsymbol{q}_1) = e^{t\Omega}\boldsymbol{q}_1$, with Ω skew-symmetric $N \times N$ (non-zero) matrix.[3] If U is invariant with respect to the above-mentioned one-parameter group of rotations $\varphi^t = e^{t\Omega}$, a relative equilibrium satisfies the equation

$$\Omega^2\boldsymbol{q} = \nabla_M U(\boldsymbol{q}) \ .$$

It follows that if $\dim(E) = 2$, then $\Omega = \begin{bmatrix} 0 & -\omega \\ \omega & 0 \end{bmatrix}$ with $\omega \in \mathbb{R}$, $\omega \neq 0$, and therefore such a relative equilibrium configuration is a central configuration, $-\omega^2\boldsymbol{q} = \nabla_M U(\boldsymbol{q})$. Conversely, planar central configurations with $U(\boldsymbol{q}) > 0$ (if U is homogeneous of negative degree) yield relative equilibria, with a suitable (angular speed) ω.

If $\dim(E) = 3$, then since the non-zero 3×3 skew-symmetric matrix Ω has rank 2, E can be written as $\ker\Omega \oplus E'$, where $\ker\Omega$ is the fixed direction of the rotations $\varphi^t = e^{t\Omega}$, and E' is the orthogonal complement of $\ker\Omega$. In a suitable reference, $\Omega = \begin{bmatrix} 0 & -\omega & 0 \\ \omega & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$. For further reference, let

$$P\colon E \to E' \tag{1.7}$$

denote the orthogonal projection.

If U is the homogeneous Newtonian potential of (1.1) with $\alpha > 0$ and $m_j > 0$, then it is easy to see that relative equilibrium configurations must belong to the plane E'. This is not true in general: it is possible to find examples of relative equilibria which are not planar – see (4.1) (and hence they are not central configurations). For more on equilibrium (and homographic) solutions: [19] (§369–§382bis at pp. 284–306), [4], [2].

Recent and non recent relevant literature on central configurations is the following: [14], [7], [16, 17], [10, 11, 12], [8], [20], [1], [6], [3].

From now on, unless otherwise stated, assume that U is invariant with respect to all isometries in E, all masses $m_j > 0$ are positive, and U is homogeneous of negative degree $-\alpha$.

The potential U is invariant with respect to a suitable subgroup of $\Sigma_n \times O(E)$, where Σ_n is the symmetric group on n elements and $O(E)$ denotes the orthogonal group on the euclidean space E. For example, if all masses are equal and U is defined as in (1.1), then $G = \Sigma_n \times O(E)$; if all masses are distinct, then $G = \{1\} \times O(E)$.

The following proposition is a well-known characterization of the set of central configurations, which we will generalize to relative equilibria in (1.9).

(1.8) *Let $S \subset Y$ denote the inertia ellipsoid, defined as $S = \{\boldsymbol{q} \in Y : \|\boldsymbol{q}\|_M^2 = 1\}$. A point $\boldsymbol{q} \in S$ is a central configuration if and only if it is a critical point of the restriction of U to S.*

[3] Cf. [18], p. 401.

Proof. Critical points of $U|_S$ are points $\boldsymbol{q} \in Y$ such that $\ker dU \supset T_{\boldsymbol{q}}S$. With respect to the (non-degenerate) bilinear form $\langle -, - \rangle_M$, this can be written as $\nabla_M U(\boldsymbol{q}) = \lambda \nabla_M (\|\boldsymbol{q}\|^2_M) = 2\lambda \boldsymbol{q}$. *q.e.d.*

(1.9) *Assume* $\dim(E) = 3$. *Let* C *be the* vertical cylinder *defined as*

$$C = \{\boldsymbol{q} \in Y : \langle P\boldsymbol{q}, \boldsymbol{q} \rangle_M = c \ , \}$$

where P *is the projection of* E *to* E' *as in (1.7) and* $c = \langle P\overline{\boldsymbol{q}}, \overline{\boldsymbol{q}} \rangle_M$. *A configuration* $\overline{\boldsymbol{q}} \in Y$ *is a relative equilibrium configuration rotating by* $e^{t\Omega}$ *if and only if it is a critical point of* U *restricted to* $C \subset X$ *and* $U(\overline{\boldsymbol{q}}) > 0$.

Proof. The configuration $\boldsymbol{q}$ is a relative equilibrium configuration if and only if $\Omega^2 \boldsymbol{q} = \nabla_M U(\boldsymbol{q})$; since $\Omega^2 = -\omega^2 P$ for an $\omega \neq 0$, this is equivalent to

$$\nabla_M U(\boldsymbol{q}) = -\omega^2 P\boldsymbol{q}.$$

On the other hand, $\nabla_M \left(\langle P\boldsymbol{q}, \boldsymbol{q} \rangle_M \right) = 2P\boldsymbol{q}$, hence $\boldsymbol{q} \in C$ is a critical point of the restriction $U|_C$ iff $\nabla_M U(\boldsymbol{q}) = 2\lambda P\boldsymbol{q}$. The proof follows since, by homogeneity,

$$\langle \nabla_M U(\boldsymbol{q}), \boldsymbol{q} \rangle_M = -\alpha U(\boldsymbol{q})$$

and $\langle P\boldsymbol{q}, \boldsymbol{q} \rangle_M = \|P\boldsymbol{q}\|^2_M$. *q.e.d.*

(1.10) *Let* K *be a subgroup of the symmetry group* G *of* U *on* Y. *Then the inertia ellipsoid* S *is* K*-invariant, and critical points of the restriction of* U *to* $S^K = \{\boldsymbol{q} \in S : K\boldsymbol{q} = \boldsymbol{q}\}$ *are precisely the critical points of* $U|_S$ *belonging to* S^K. *If the vertical cylinder* C *is* K*-invariant, then critical points of the restriction of* U *to* $C^K = \{\boldsymbol{q} \in C : K\boldsymbol{q} = \boldsymbol{q}\}$ *are precisely the critical points of* $U|_C$ *belonging to* C^K.

Proof. The Palais principle of Symmetric Criticality [9] states that *in order for a symmetric point* p *to be a critical point it suffices that it be a critical point of* $f|_\Sigma$, *the restriction of* f *to* Σ, where $f \colon M \to \mathbb{R}$ is a smooth K-invariant function on M, M is a smoorh manifold and a symmetric point is an element of $\Sigma = M^K = \{p \in M : gp = p \text{ for all } g \in K\}$. Thus, the statement follows by setting $f = U|_M$ with $M = S$ or $M = C$. *q.e.d.*

2 Central configurations as (equivariant) fixed points.

In [5] a way to relate planar central configurations to projective classes of fixed points was introduced. We now generalize the results therein to arbitrary dimensions.

Consider a homogeneous potential U, as above with the further assumption that $\forall \boldsymbol{q}, U(\boldsymbol{q}) > 0$. From this it follows that $\nabla_M U(\boldsymbol{q}) \neq \mathbf{0}$ because $\langle \nabla_M U(\boldsymbol{q}), \boldsymbol{q} \rangle_M = -\alpha U(\boldsymbol{q})$.

(2.1) *The map $F\colon S \to S$ defined as*

$$F(\boldsymbol{q}) = -\frac{\nabla_M U(\boldsymbol{q})}{\|\nabla_M U(\boldsymbol{q})\|_M}$$

is well-defined, and a configuration $\bar{\boldsymbol{q}} \in S$ is a central configuration if and only if it is a fixed point of F.

Proof. It follows from the assumption that $\forall \boldsymbol{q}, \nabla_M U(\boldsymbol{q}) \neq \mathbf{0}$, and therefore F is well-defined. A configuration $\boldsymbol{q}$ is fixed by F if and only if there exists $\lambda = \|\nabla_M U(\boldsymbol{q})\|_M > 0$ such that $\nabla_M U(\boldsymbol{q}) = -\lambda \boldsymbol{q}$. Hence, if $F(\boldsymbol{q}) = \boldsymbol{q}$ then $\boldsymbol{q}$ is central. Conversely, by homogeneity of U, $\langle \nabla_M U(\boldsymbol{q}), \boldsymbol{q} \rangle_M = -\alpha U(\boldsymbol{q})$, and hence if $\boldsymbol{q} \in S$ is a central configuration then $\nabla_M U(\boldsymbol{q}) = \lambda \boldsymbol{q} \implies \lambda = -\alpha U(\boldsymbol{q}) < 0$ and therefore $F(\boldsymbol{q}) = \boldsymbol{q}$. *q.e.d.*

Now, in general (and without the positivity assumption) the map F needs not being compactly fixed (see for example Robert's continuum [13] of central configurations with four unit masses and a fifth negative $-1/4$ mass in the origin[4]). In the gravitational case (positive masses and Newtonian mutual attraction), the map F turns out to be compactly fixed [5] (see also Shub's estimates [15]).

(2.2) *Let G be the symmetry group of U on Y, as above. Then F is G-equivariant (that is, for each $g \in G$ and for each $\boldsymbol{q} \in S$, $F(g\boldsymbol{q}) = gF(\boldsymbol{q})$). For each subgroup $K \subset G$, F induces a self-map $\bar{F}\colon S/K \to S/K$ on the quotient space S/K.*

Proof. For $g \in G$, $gS = S$, and for each g such that $U(g\boldsymbol{q}) = U(\boldsymbol{q})$ the equality $\nabla_M U(g\boldsymbol{q}) = g\nabla_M U(\boldsymbol{q})$ holds. In fact, since $U \circ g = U$, $dU \circ g = dU$, and therefore for each vector $\boldsymbol{v}$ one has $\langle \nabla_M U(\boldsymbol{q}), \boldsymbol{v} \rangle_M = dU(\boldsymbol{q})[\boldsymbol{v}] = dU(g\boldsymbol{q})[g\boldsymbol{v}] = \langle \nabla_M U(g\boldsymbol{q}), g\boldsymbol{v} \rangle_M = \langle g^{-1}\nabla_M U(g\boldsymbol{q}), \boldsymbol{v} \rangle_M$. Thus F is G-equivariant, and hence K-equivariant for each subgroup $K \subset G$. *q.e.d.*

Let U be a homogeneous potential with the following property: for each orthogonal projection $p\colon E \to \eta$ on a plane η, for each $\boldsymbol{q} \in S$ there exists $j \in \{1, \dots, n\}$ such that

$$p\left(\frac{\partial U}{\partial \boldsymbol{q}_j}(\boldsymbol{q})\right) \cdot p(\boldsymbol{q}_j) \leq 0 \ . \tag{2.3}$$

Moreover, if there exists $i \in \{1, \dots, n\}$ such that $p(\boldsymbol{q}_i) \neq 0$, then the j of (2.3) is such that $p(\boldsymbol{q}_j) \neq 0$.

[4] In the AMS review of [13], D. Saari states that "a similar effect occurs for positive masses, but with two or more homogeneous potentials". More precisely (in review MR1695029), "should the potential be a sum of homogeneous potentials, then there always is a continuum of different relative equilibria configurations". In review MR2245515: "there can be a continuum of different central configurations parametrized by the value of the moment of inertia".

It is easy to see that the Newtonian potential (1.1) (with positive masses and homogeneity $-\alpha$) satisfies Property (2.3): let j be the index maximizing $\|p(\boldsymbol{q}_k)\|^2$ for $k = 1, \dots, n$; then

$$\begin{aligned} p\left(\alpha \sum_{k\neq j} \frac{m_j m_k(\boldsymbol{q}_k - \boldsymbol{q}_j)}{\|\boldsymbol{q}_k - \boldsymbol{q}_j\|^{\alpha+2}}\right) \cdot p(\boldsymbol{q}_j) &= \alpha \left(\sum_{k\neq j} \frac{m_j m_k(p(\boldsymbol{q}_k) - p(\boldsymbol{q}_j))}{\|\boldsymbol{q}_k - \boldsymbol{q}_j\|^{\alpha+2}}\right) \cdot p(\boldsymbol{q}_j) \\ &= \alpha \sum_{k\neq j} \frac{m_j m_k(p(\boldsymbol{q}_k)\cdot p(\boldsymbol{q}_j) - \|p(\boldsymbol{q}_j)\|^2)}{\|\boldsymbol{q}_k - \boldsymbol{q}_j\|^{\alpha+2}} \\ &\leq \alpha \sum_{k\neq j} \frac{m_j m_k(\|p(\boldsymbol{q}_k)\|\|p(\boldsymbol{q}_j)\| - \|p(\boldsymbol{q}_j)\|^2)}{\|\boldsymbol{q}_k - \boldsymbol{q}_j\|^{\alpha+2}} \\ &\leq 0 \end{aligned}$$

It is trivial to see that if U satisfies (2.3), then the map $F\colon S \to S$ defined in (2.1) satisfies the following property: for each orthogonal projection $p\colon E \to P$ on a plane P, for each $\boldsymbol{q} \in S$, there exists $j \in \{1, \dots, n\}$ such that

$$p(F_j(\boldsymbol{q})) \cdot p(\boldsymbol{q}_j) \geq 0 \ . \tag{2.4}$$

(2.5) Theorem. *Suppose that $G \supset SO(E)$, where as above G is the symmetry group of U on Y, and $SO(E) = SO(d)$ (the group of rotations in E) acting diagonally on Y. Let $\overline{F}\colon S/SO(d) \to S/SO(d)$ be the map induced on the quotient space by (2.2). Assume that all masses are positive, $U > 0$ and (2.4) holds. Then, if $\pi\colon S \to S/SO(d)$ denotes the projection on the quotient,*

$$\pi(\mathrm{Fix}(F)) = \mathrm{Fix}(\overline{F}) \ .$$

Proof. If $\boldsymbol{q} \in \mathrm{Fix}(F) \subset S$ is fixed by F, then $\overline{F}(\pi(\boldsymbol{q})) = \pi(F(\boldsymbol{q})) = \pi(\boldsymbol{q}) \implies \pi(\boldsymbol{q}) \in \mathrm{Fix}(\overline{F})$, hence $\pi(\mathrm{Fix}(F)) \subset \mathrm{Fix}(\overline{F})$. On the other hand, let $\pi(\boldsymbol{q}) \in \mathrm{Fix}(\overline{F})$. Then there exists a rotation $g \in SO(E)$ such that $F(\boldsymbol{q}) = g\boldsymbol{q}$. Without loss of generality, we can assume that $g = e^{\Omega}$, where Ω is an antisymmetric $d \times d$ matrix, with k (2×2)-blocks on the diagonal $\begin{bmatrix} 0 & -\theta_i \\ \theta_i & 0 \end{bmatrix}$, with $\theta_i \in [-\pi, \pi]$ for each $i = 1, \dots, k$, and, if d is odd, a one-dimensional diagonal zero entry ($d = 2k$ or $d = 2k+1$). We can also assume that only the first (say, $l \leq k$) blocks have $\theta_i \neq 0$, hence Ω has l non-zero (2×2) diagonal blocks and is zero outside. Note that for each $\boldsymbol{x} \in E$, the quadratic form $(e^{\Omega}\boldsymbol{x}) \cdot (\Omega\boldsymbol{x})$ on E can be written with l non-singular positive-defined blocks

$$\theta_i \begin{bmatrix} \sin\theta_i & -\cos\theta_i \\ \cos\theta_i & \sin\theta_i \end{bmatrix} \sim \theta_i \begin{bmatrix} \sin\theta_i & 0 \\ 0 & \sin\theta_i \end{bmatrix}$$

on the diagonal, and hence it is non-negative. Here the symbol "$\sim$" means that the two matrices yield the same quadratic form. Moreover, if one writes

$\boldsymbol{x} \in E \cong \mathbb{R}^d$ as $(\boldsymbol{z}_1, \boldsymbol{z}_2, \dots, \boldsymbol{z}_l, x_{2l+1}, \dots, x_d)$, with $\boldsymbol{z}_i \in \mathbb{R}^2$ for $i = 1, \dots, l$ and $x_i \in \mathbb{R}$, then

$$(e^{\Omega}\boldsymbol{x}) \cdot (\Omega\boldsymbol{x}) = \sum_{i=1}^{l} \theta_i \sin\theta_i \|\boldsymbol{z}_i\|^2 \ . \tag{2.6}$$

Let $p_i \colon E \to \mathbb{R}^2$ denote the projection $\boldsymbol{x} \mapsto \boldsymbol{z}_i$, for $i = 1, \dots, l$.

Since $F(e^{t\Omega}\boldsymbol{q})$ does not depend on $t \in \mathbb{R}$,

$$\begin{aligned} 0 &= \frac{d}{dt}\|F(e^{t\Omega}\boldsymbol{q})\|_M^2|_{t=0} \\ &= 2\langle F(\boldsymbol{q}), \Omega\boldsymbol{q}\rangle_M \\ &= 2\langle e^{\Omega}\boldsymbol{q}, \Omega\boldsymbol{q}\rangle_M \\ &= 2\sum_{j=1}^{n} m_j (e^{\Omega}\boldsymbol{q}_j) \cdot (\Omega\boldsymbol{q}_j) \end{aligned}$$

For each $j = 1, \dots, n$ the inequality $m_j > 0$ holds, and for each $\boldsymbol{x} \in E$ the inequality $(e^{\Omega}\boldsymbol{x}) \cdot (\Omega\boldsymbol{x}) \geq 0$ holds: it follows that for each j, $(e^{\Omega}\boldsymbol{q}_j) \cdot (\Omega\boldsymbol{q}_j) = 0$. By (2.6), this implies that, given j, for each $i = 1, \dots, l$ either $p_i(\boldsymbol{q}_j) = 0$ or $\theta_i \in \{\pi, -\pi\}$ (since $\theta_i \neq 0$ for $i = 1, \dots, l$). If $p_i(\boldsymbol{q}_j) = 0$ for each j, then actually $g\boldsymbol{q} = \boldsymbol{q}$, and therefore $\pi(\boldsymbol{q}) \in \pi(\mathrm{Fix}(F))$. So, without loss of generality one can assume that for each $i = 1, \dots, l$ there exists j such that $p_i(\boldsymbol{q}_j) \neq 0$. Suppose that $l \geq 1$, and therefore $\theta_1 = \pm\pi$. By property (2.4) there exists $\bar{j}$ such that $p_1(F_{\bar{j}}(\boldsymbol{q})) \cdot p_1(\boldsymbol{q}_{\bar{j}}) \geq 0$ and $p_1(\boldsymbol{q}_{\bar{j}}) \neq 0$. But this would imply that

$$F(\boldsymbol{q}) = g\boldsymbol{q} = e^{\Omega}\boldsymbol{q} \implies -p_1(\boldsymbol{q}_{\bar{j}}) \cdot p_1(\boldsymbol{q}_{\bar{j}}) \geq 0 \implies p_1(\boldsymbol{q}_{\bar{j}}) = 0 \ ,$$

which is not possible. Therefore, if condition (2.4) holds, $l = 0$, and $g\boldsymbol{q} = \boldsymbol{q}$. The conclusion follows. *q.e.d.*

3 Projective fixed points and Morse indices

In this section, we finally prove the equation relating fixed point and Morse indices of central configurations.

(3.1) *If U is homogeneous of degree $-\alpha$, then for each central configuration $\boldsymbol{q}$, up to a linear change of coordinates*

$$-\alpha U(\boldsymbol{q})(I - F'(\boldsymbol{q})) = D^2\tilde{U}(\boldsymbol{q}),$$

where $F \colon S \to S$ is the function of (2.1), defined as $F(\boldsymbol{q}) = -\dfrac{\nabla_M U(\boldsymbol{q})}{\|\nabla_M U(\boldsymbol{q})\|_M}$, and $D^2\tilde{U}(\boldsymbol{q})$ is the Hessian of the restriction $\tilde{U}$ of U to S, evaluated at $\boldsymbol{q}$.

Proof. As above, let $X = E^n \smallsetminus \Delta$ denote the configuration space of E. After a linear change of coordinates in X, we can assume $m_i = 1$ for each i and $\boldsymbol{q} = (1, 0, \dots, 0) = (1, \mathbf{0})$ (rescale $\boldsymbol{q}$ by a diagonal matrix with suitable entries on its diagonal, and apply a rotation - this leaves U homogeneous of the same degree). Given suitable linear coordinates $\boldsymbol{x} = (x_0, x_1, \dots, x_l)$ in $Y \cong \mathbb{R}^{l+1}$, the ellipsoid S has equation $\|\boldsymbol{x}\|^2 = 1$, and $F(\boldsymbol{x}) = -\dfrac{\nabla U}{\|\nabla U\|}$, in the $\boldsymbol{x}$-coordinates. Therefore, if $\boldsymbol{u} = (u_1, \dots, u_l) \mapsto (\sqrt{1 - \|\boldsymbol{u}\|^2}, u_1, \dots, u_l) \in S$ is a local chart around the central configuration $\boldsymbol{q} \sim (1, \mathbf{0})$,

$$\begin{aligned} \frac{\partial F_\alpha}{\partial u_\beta}(\mathbf{0}) &= \frac{\partial^2 U}{\partial x_\alpha \partial x_\beta}(\boldsymbol{q}) \|\nabla U(\boldsymbol{q})\|^{-1} \\ D^2_{\alpha\beta}\tilde{U}(\mathbf{0}) &= \frac{\partial^2 U}{\partial x_\alpha \partial x_\beta}(\boldsymbol{q}) - \delta_{\alpha\beta}\frac{\partial U}{\partial x_0}(\boldsymbol{q}) \ . \end{aligned} \tag{3.2}$$

Now, $\dfrac{\partial U}{\partial x_0} = \langle \nabla U, \boldsymbol{q} \rangle = -\alpha U(\boldsymbol{q})$, and since $\boldsymbol{q}$ is a central configuration, it is a fixed point of F and therefore

$$F(\boldsymbol{q}) = \boldsymbol{q} \implies \frac{\partial U}{\partial x_\alpha} = 0, \quad \text{for } \alpha = 1, \dots, l$$

and

$$\frac{\partial U}{\partial x_0}(\boldsymbol{q}) = -\|\nabla U(\boldsymbol{q})\| = -\alpha U(\boldsymbol{q}) \ . \tag{3.3}$$

From (3.3) and (3.2) it follows that for $\alpha, \beta = 1, \dots, l$,

$$\begin{aligned} D^2_{\alpha\beta}\tilde{U}(\mathbf{0}) + \delta_{\alpha\beta}\frac{\partial U}{\partial x_0}(\boldsymbol{q}) = \frac{\partial^2 U}{\partial x_\alpha \partial x_\beta}(\boldsymbol{q}) = \frac{\partial F_\alpha}{\partial u_\beta}(\mathbf{0}) \|\nabla U(\boldsymbol{q})\| \\ \implies D^2\tilde{U}(\mathbf{0}) = -\alpha U(\boldsymbol{q}) I + \alpha U(\boldsymbol{q}) F'(\boldsymbol{q}) \end{aligned} \tag{3.4}$$

q.e.d.

The *maximal isotropy stratum* in $S/SO(d)$ is the set of all orbits of points in S with trivial isotropy (and hence with maximal isotropy type). It is an open (and dense) subspace of $S/SO(d)$.

(3.5) Corollary. *Assume the hypotheses of Theorem (2.5) hold. Then for each non-degenerate projective class of central configurations $\boldsymbol{q} \in \mathrm{Fix}(\bar{F})$ in the maximal isotropy stratum of $\bar{S} = S/SO(d)$, the fixed point index $\mathrm{ind}(\boldsymbol{q}, \bar{F})$ and the Morse index $\mu_{\tilde{U}}(\boldsymbol{q})$ of $\tilde{U}$ at $\boldsymbol{q}$ are related by the equality*

$$\mathrm{ind}(\boldsymbol{q}, \bar{F}) = (-1)^{\mu_{\tilde{U}}(\boldsymbol{q}) + \epsilon},$$

where $\epsilon = d(n-1) - 1 - d(d-1)/2$.

Proof. Since $\tilde{U}$ is $SO(d)$-invariant (with diagonal action), and the $SO(d)$-orbits in S with maximal isotropy type have dimension $d(d-1)/2$, the Hessian $D^2\tilde{U}$ has $d(d-1)/2$-dimensional kernel, if $\boldsymbol{q}$ is non-degenerate. By proposition (3.1), if follows that F' has a $d(d-1)/2$-multiple eigenvalue 1, and $\mathrm{ind}(\boldsymbol{q}, \bar{F}) = (-1)^c$, where c is the number of negative eigenvalues of $-D^2\tilde{U}$, i.e. the number of positive eigenvalues of $D^2\tilde{U}$, which is equal to $\dim(S) - d(d-1)/2 - \mu_{\tilde{U}}(\boldsymbol{q})$. Therefore

$$\mathrm{ind}(\boldsymbol{q}, \bar{F}) = (-1)^{\dim(S)-d(d-1)/2-\mu_{\tilde{U}}(\boldsymbol{q})} = (-1)^{\epsilon-\mu_{\tilde{U}}(\boldsymbol{q})} = (-1)^{\mu_{\tilde{U}}(\boldsymbol{q})+\epsilon}$$

where $\epsilon = \dim(S) - d(d-1)/2 = d(n-1) - 1 - d(d-1)/2$. *q.e.d.*

(3.6) Corollary. *Let U be the Newton potential in* (1.1)*, with positive masses, $\alpha > 0$, and $\dim(E) = 2$. Then $X \approx \mathbb{C}^n \setminus \Delta$, and $S/SO(2) \approx \mathbb{P}^{n-2}(\mathbb{C})_0 \subset \mathbb{P}^{n-2}(\mathbb{C})$, where $\mathbb{P}^{n-2}(\mathbb{C})_0$ is the subset of $\mathbb{P}^{n-1}(\mathbb{C})$ defined in projective coordinates as $\mathbb{P}^{n-2}(\mathbb{C})_0 \cong \{[z_1 : \ldots, z_n] \in \mathbb{P}^n : \sum_j m_j z_j = 0, \forall i, j, z_i \neq z_j\}$. Then for each non-degenerate projective class of central configurations $q \in \mathrm{Fix}(\overline{F})$, with $\overline{F}\colon \mathbb{P}_0^{n-2}(\mathbb{C}) \to \mathbb{P}^{n-2}(\mathbb{C})$,*

$$\mathrm{ind}(\boldsymbol{q}, \overline{F}) = (-1)^{\mu_{\overline{U}}(\boldsymbol{q})} \ .$$

Proof. If $d = 2$, then $d(n-1) - 1 - d(d-1)/2 = 2(n-2)$. *q.e.d.*

4 An example of non-planar relative equilibrium

(4.1) Example (Non-central and non-planar equilibrium solution)**.** In $E \cong \mathbb{R}^3$, let R_x, R_y and R_z denote rotations of angle π around the three coordinate axes. Fix three non-zero constants c_1, c_2, c_3. Consider the problem with 6 bodies in E, symmetric with respect to the group K with non-trivial elements of $\Sigma_6 \times SO(3)$

$$((34)(56), R_x), ((12)(56), R_y), ((12)(34), R_z) \ .$$

Assume $m_j = 1$, for $j = 1, \ldots, 6$, and let U be the potential defined on E^6 by

$$U(\boldsymbol{q}) = \sum_{i<j} \frac{1 - \gamma_i\gamma_j}{\|\boldsymbol{q}_i - \boldsymbol{q}_j\|}$$

where

$$\gamma_1 = \gamma_2 = c_1, \quad \gamma_3 = \gamma_4 = c_2, \quad \gamma_5 = \gamma_6 = c_3 \ .$$

Now, U is invariant with respect to K, and the vertical cylinder is K-invariant: it follows from (1.9) that critical points of the restriction of U to C^K are equilibrium configurations.

In other words, three pairs of bodies of unit masses, each pair charged with charge c_j, are constrained each pair to belong to one of the coordinate axes and to be symmetric with respect to the origin.

The space $X^K = Y^K$ has dimension 3, and can be parametrized by (x, y, z), where x, y and z are (respectively) the coordinates along the corresponding axis of particles 1, 3 and 5. The generic configuration $\boldsymbol{q} \in X^K$ can be written as

$$\begin{aligned} \boldsymbol{q}_1 &= (x, 0, 0) \qquad \boldsymbol{q}_2 = (-x, 0, 0) \\ \boldsymbol{q}_3 &= (0, y, 0) \qquad \boldsymbol{q}_4 = (0, -y, 0) \\ \boldsymbol{q}_5 &= (0, 0, z) \qquad \boldsymbol{q}_6 = (0, 0, -z), \end{aligned}$$

and the potential U restricted to X^K in such coordinates is

$$U(x,y,z) = \frac{1-c_1^2}{2|x|} + \frac{1-c_2^2}{2|y|} + \frac{1-c_3^2}{2|z|} + 4\frac{1-c_1c_2}{\sqrt{x^2+y^2}} + 4\frac{1-c_1c_3}{\sqrt{x^2+z^2}} + 4\frac{1-c_2c_3}{\sqrt{y^2+z^2}}\,.$$

The vertical cylinder $C^K \subset X^K$ is

$$\begin{aligned} C^K &= \{\boldsymbol{q} \in X^K : \langle P\boldsymbol{q}, \boldsymbol{q}\rangle_M = 2\} \\ &= \{(x,y,z) \in X^K : 2x^2 + 2y^2 = 2\}\,. \end{aligned}$$

Hence an equilibrium solution is a critical point of $U(\cos t, \sin t, z)$ with positive value $U > 0$. Now, assume

$$c_1 > 1, \quad c_2 < -1, \quad c_3 < -1\,. \tag{4.2}$$

Then $1 - c_1^2 < 0$, $1 - c_2^2 < 0$, $1 - c_3^2 < 0$, $1 - c_2c_3 < 0$; moreover, $1 - c_1c_2 > 0$ and $1 - c_1c_3 > 0$. The restricted potential U is defined as

$$U(t,z) = \frac{1-c_1^2}{2|\cos t|} + \frac{1-c_2^2}{2|\sin t|} + \frac{1-c_3^2}{2z} + 4(1-c_1c_2) + 4\frac{1-c_1c_3}{\sqrt{\cos^2 t + z^2}} + 4\frac{1-c_2c_3}{\sqrt{\sin^2 t + z^2}},$$

and is defined on the strip $(t, z) \in T = (0, \pi/2) \times (0, +\infty)$.

If

$$1 - c_1^2 + 8(1 - c_1c_3) < 0, \tag{4.3}$$

then for each (t, z)

$$\begin{aligned} \frac{1-c_1^2}{2|\cos t|} + 4\frac{1-c_1c_3}{\sqrt{\cos^2 t + z^2}} &< \frac{1-c_1^2}{2|\cos t|} + 4\frac{1-c_1c_3}{|\cos t|} \\ &= \frac{1}{2|\cos t|}\left(1 - c_1^2 + 8(1-c_1c_3)\right) < 0 \end{aligned}$$

and hence $U(t, z) < 4(1 - c_1c_2)$ for each $(t, z) \in T$ and $U \to -\infty$ on the boundary of T.

Furthermore, for each $t \in (0, \pi/2)$,

$$\begin{aligned} \frac{\partial U}{\partial z} &= \frac{c_3^2 - 1}{2z^2} + 4z\frac{c_2c_3 - 1}{(z^2 + \sin^2 t)^{3/2}} - 4z\frac{1-c_1c_3}{(z^2+\cos^2 t)^{3/2}} \\ &< \frac{c_3^2-1}{2z^2} + 4\frac{c_2c_3-1}{z^2} - 4z\frac{1-c_1c_3}{(z^2+1)^{3/2}}\,. \end{aligned}$$

It follows that if

$$c_3^2 - 1 + 8(c_2c_3 - 1) < 8(1 - c_1c_3), \tag{4.4}$$

then for every $t \in (0, \pi/2)$ there exists z_0 such that $z > z_0 \implies \frac{\partial U}{\partial z}(t, z) < 0$.

Thus, whenever both (4.3) and (4.4) hold, the restriction $U(t, z)$ attains its maximum in the interior of the strip T. Such a maximum (t_m, z_m) corresponds to an equilibrium configuration if and only if $U(t_m, z_m) > 0$, from (1.9). Note that if $t = \pi/4$ and $z = 1/2$ then

$$U(\pi/4, z) = \frac{\sqrt{2}}{2}(2 - c_1^2 - c_2^2) + 4(1 - c_1c_2) + 1 - c_3^2 + \frac{4(2 - c_3(c_1 + c_2))}{\sqrt{3/4}},$$

which is positive, for example, if $c_1 = 20$ and $c_2 = c_3 = -2$. Such coefficients satisfy (4.2), (4.3) and (4.4), and therefore for such a choice of c_i there exist non-planar relative equilibrium configurations.

References

[1] Albouy, A. Symétrie des configurations centrales de quatre corps. *C. R. Acad. Sci. Paris Sér. I Math. 320*, 2 (1995), 217–220.

[2] Albouy, A., and Chenciner, A. Le problème des n corps et les distances mutuelles. *Invent. Math. 131*, 1 (1998), 151–184.

[3] Albouy, A., and Kaloshin, V. Finiteness of central configurations of five bodies in the plane. *Ann. of Math. (2) 176*, 1 (2012), 535–588.

[4] Betti, E. Sopra il moto di un sistema di un numero qualunque di punti che si attraggono o si respingono tra loro. *Annali di Matematica Pura ed Applicata (1867-1897) 8*, 1 (1877), 301–311.

[5] Ferrario, D. L. Planar central configurations as fixed points. *J. Fixed Point Theory Appl. 2*, 2 (2007), 277–291.

[6] Hampton, M., and Moeckel, R. Finiteness of relative equilibria of the four-body problem. *Invent. Math. 163*, 2 (2006), 289–312.

[7] Moeckel, R. On central configurations. *Math. Z. 205*, 4 (1990), 499–517.

[8] Pacella, F. Central configurations of the N-body problem via equivariant Morse theory. *Arch. Rational Mech. Anal. 97*, 1 (1987), 59–74.

[9] Palais, R. S. The principle of symmetric criticality. *Comm. Math. Phys. 69*, 1 (1979), 19–30.

[10] Palmore, J. I. Classifying relative equilibria. I. *Bull. Amer. Math. Soc. 79* (1973), 904–908.

[11] Palmore, J. I. Classifying relative equilibria. II. *Bull. Amer. Math. Soc. 81* (1975), 489–491.

[12] Palmore, J. I. Classifying relative equilibria. III. *Lett. Math. Phys. 1*, 1 (1975/76), 71–73.

[13] Roberts, G. E. A continuum of relative equilibria in the five-body problem. *Phys. D 127*, 3-4 (1999), 141–145.

[14] Saari, D. G. *Collisions, rings, and other Newtonian N-body problems*, vol. 104 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2005.

[15] Shub, M. Appendix to Smale's paper: "Diagonals and relative equilibria". In *Manifolds – Amsterdam 1970 (Proc. Nuffic Summer School)*, Lecture Notes in Mathematics, Vol. 197. Springer, Berlin, 1971, pp. 199–201.

[16] Smale, S. Topology and mechanics. I. *Invent. Math. 10* (1970), 305–331.

[17] Smale, S. Topology and mechanics. II. The planar n-body problem. *Invent. Math. 11* (1970), 45–64.

[18] Spivak, M. *A comprehensive introduction to differential geometry. Vol. I*, second ed. Publish or Perish, Inc., Wilmington, Del., 1979.

[19] Wintner, A. *The Analytical Foundations of Celestial Mechanics*. Princeton Mathematical Series, v. 5. Princeton University Press, Princeton, N. J., 1941.

[20] Xia, Z. Central configurations with many small masses. *J. Differential Equations 91*, 1 (1991), 168–179.